

Organic Feeding Schedule (10 -week Bloom)



“Moderate” Program

USEFUL CONVERSIONS

1 teaspoon	=	5 ml
1 Tablespoon	=	15 ml
1 ounce	=	30 ml
1 quart	=	946 ml
1 gallon	=	3.785 L
1 gallon	=	128 oz
*1 teaspoon (powder)	=	2 1/3 grams (approx.)

	Grow Week 1	Grow Week 2	Grow Week 3	Grow Week 4	Bloom Week 1	Bloom Week 2	Bloom Week 3	Bloom Week 4	Bloom Week 5	Bloom Week 6	Bloom Week 7	Bloom Week 8	Bloom Week 9	Bloom Week 10
GROW NATURAL	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	4ml ▶▶ gal	4ml ▶▶ gal	4ml ▶▶ gal	3ml ▶▶ gal	Flush
BLOOM NATURAL					4ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	7ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	12ml ▶▶ gal	10ml ▶▶ gal	Flush
DEUCEDEUCE						1ml ▶▶ gal	1ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	3ml ▶▶ gal	3ml ▶▶ gal	4ml ▶▶ gal	4ml ▶▶ gal	Flush
MAYAN MICROZYME	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal							Flush
FLAVORFUL	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal					Flush
HUM-BOLT	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	3ml ▶▶ gal	3ml ▶▶ gal	4ml ▶▶ gal	4ml ▶▶ gal	Flush
HUMBOLDT HONEY ES	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	15ml ▶▶ gal
HUMBOLDT ROOTS	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	1ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal							Flush
PROZYME	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	2ml ▶▶ gal	5ml ▶▶ gal	5ml ▶▶ gal	7ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	10ml ▶▶ gal	Flush
MYCOMADNESS	1/4tsp ▶▶	1/4tsp ▶▶	1/4tsp ▶▶	1/4tsp ▶▶	1/2sp ▶▶ gal	1/2Tsp ▶▶	1/2tsp ▶▶							Flush
GEOLIFE	1st Application	2nd Application			3rd Application									
MYCOMAXIMUM	Add to soil prior to planting : Ratio is 4 oz. per 1.5 cubic feet or 2 tbsp. per gallon container.													Flush

Always use un-chlorinated water, maintain pH levels between 5.5 -7.2 and check reservoir after adding all nutrients. Oxygenate water before and during application. To prevent nutrient settling, always use a pump at the bottom of the reservoir to continually agitate and mix the nutrient water during application. Research and Development conducted using water obtained by reverse osmosis containing near 0 PPM. Humboldt Nutrients organic feeding schedules work best in soil applications.

For optimum biological activity which fixates and increases availability of nutrients, combine nutrients and additives with water and allow to “brew” for at least 24 hours prior to use.

